

Coronavirus (COVID 19) Risk Assessment		
Establishment: Roundabout pre-School	Assessment by: Adele Laycock	Date: 21/9/20
Review date: As and when the government guidance changes 2020	Red- High Yellow - medium risk Green- little or no risk	

Amendment 1

Focus	Area of consideration	Recommendation	Risks and level of risks
Children	If a child/adult starts to display symptoms	<ul style="list-style-type: none"> If anyone in the setting has symptoms of coronavirus (COVID-19): a high temperature, new and persistent cough or a loss of, or change in, normal sense of taste or smell (anosmia), however mild, they should self-isolate for at least 10 days from when their symptoms started; or if they are not experiencing symptoms, but have tested positive for coronavirus (COVID-19), they should self-isolate for at least 10 days starting from the day the test was taken. If a child is awaiting collection, they should be moved, to an area where they can be isolated with appropriate adult supervision (front of the building use the bench). If they need to go to the bathroom while waiting to be collected, they will be taken to a separate bathroom. The bathroom will then be cleaned and disinfected using standard cleaning products before being used by anyone else. 	

		<ul style="list-style-type: none">• PPE must be worn by staff caring for the child while they await collection if a distance of 2 metres cannot be maintained (such as for a very young child or a child with complex needs).• Any members of staff who have helped someone with symptoms and any children who have been in close contact with them do not need to go home to self-isolate. However, they must self-isolate and arrange for a test if they develop symptoms themselves (in which case, they should arrange a test), if the symptomatic person subsequently tests positive (see below) or they have been requested to do so by NHS Test and Trace.• Everyone must wash their hands thoroughly for 20 seconds with soap and running water or use hand sanitiser after any contact with someone who is unwell. The area around the person with symptoms must be cleaned with normal household disinfectant after they have left to reduce the risk of passing the infection on to other people.• If clinical advice is needed, the setting staff, parent or guardian should go online to NHS 111 (or call 111 if they don't have internet access).	
--	--	--	--